

THE MOMENT PROBLEM WITH BOUNDED DENSITY

JEAN B. LASSERRE

ABSTRACT. Let μ be a given Borel measure on $\mathbf{K} \subseteq \mathbb{R}^n$ and let $y = (y_\alpha)$, $\alpha \in \mathbb{N}^n$, be a given sequence. We provide several conditions linking y and the moment sequence $z = (z_\alpha)$ of μ , for y to be the moment sequence of a Borel measure ν on $\mathbf{K}$ which is absolutely continuous with respect to μ and such that its density is in $L_\infty(\mathbf{K}, \mu)$.

The conditions are necessary and sufficient if $\mathbf{K}$ is a compact basic semi-algebraic set, and sufficient if $\mathbf{K} \equiv \mathbb{R}^n$. Moreover, arbitrary finitely many of these conditions can be checked by solving either a semidefinite program or a linear program with a single variable!

1. INTRODUCTION

We consider the following moment problem: Let μ be a given Borel measure μ on $\mathbf{K} \subseteq \mathbb{R}^n$ and let $y = (y_\alpha)$, $\alpha \in \mathbb{N}^n$, be a given sequence. Under what conditions on y , do we have

$$(1) \quad y_\alpha = \int_{\mathbf{K}} X^\alpha h d\mu, \quad \forall \alpha \in \mathbb{N}^n,$$

for some $0 \leq h \in L_\infty(\mathbf{K}, \mu)$? (The notation X^α stands for the monomial $\prod_i X_i^{\alpha_i}$.) In Knill [6], the measure $d\nu := h d\mu$ is said to be *uniformly absolutely continuous* with respect to μ .

This moment problem (1) *with density*, initially studied by Hausdorff for the interval $[0, 1]$ of the real line, is a particular case of the general $\mathbf{K}$ -moment problem where for a given sequence y , one only asks for existence of some Borel representing measure ν on $\mathbf{K}$ (not necessarily with a density with respect to some specified measure μ). For more details on the general $\mathbf{K}$ -moment problem, the interested reader is referred to e.g. Akhiezer [1], Beg [2], Curto and Fialkow [3], Shohat and Tamarkin [12], Simon [13] and many references therein.

The moment problem *with density* is useful in e.g. probability theory, mathematical physics, statistical mechanics, potential theory and dynamical systems, and an interesting discussion on its potential applications can be found in e.g. Knill [6].

1991 *Mathematics Subject Classification.* 44A60; 90C22.

Key words and phrases. Real analysis; the moment problem; real algebraic geometry; Positivstellensatz.

Contribution. Our contribution is twofold as we consider the case where $\mathbf{K} \subset \mathbb{R}^n$ is a compact basic semi-algebraic set and the case where $\mathbf{K} \equiv \mathbb{R}^n$.

- In the former case we provide three sets of *necessary and sufficient conditions* relating y with the moment sequence z of μ . The first two sets of conditions are stated in terms of *positive definiteness* of so-called moment and localizing matrices, with entries linear in y and z , whereas the third set of conditions is stated in terms of *linear* inequalities linking the moments of μ and ν . The first two sets of (positive semidefiniteness) conditions are established thanks to powerful moment results which are the dual sides of Schmüdgen's Positivstellensatz [11], and its Putinar [10] and Jacobi and Prestel [5] refinement. Checking arbitrary finitely many of these conditions reduces to solving a *semidefinite program* with only one variable! A semidefinite program is a convex optimization problem for which efficient public softwares are now available. For more details on the theory and applications of semidefinite programming, the interested reader is referred to e.g. Vandenberghe and Boyd [14].

The third set of (linear) conditions is established thanks to a result by Vasilescu [15] (see also Krivine [7, 8]) which generalizes to basic semi-algebraic sets a previous result of Handelman [4] for polytopes (the latter being itself a generalization to polytopes of the famous Hausdorff moment conditions on the unit box $[0, 1]^n$). Checking (arbitrary) finitely many of these conditions reduces to checking whether some interval of the real line is empty.

This third set of linear conditions generalizes to compact basic semi-algebraic sets a result of Knill [6] obtained for the unit box $[0, 1]^n$. This latter result [6] states that there is some scalar $M > 0$ such that the classical Hausdorff linear moment conditions (for a measure to be supported on $[0, 1]^n$) applied to the moment sequence Mz (for some M), *dominate* those applied to y .

- In the non compact case $\mathbf{K} \equiv \mathbb{R}^n$, we provide a set of *sufficient* conditions linking y and z and assuming that z satisfies the multivariate Carleman condition. They simply state the existence of a scalar κ such that

$$0 \leq L_y(f^2) \leq \kappa \int f^2 d\mu, \quad \forall f \in \mathbb{R}[X],$$

where $L_y : \mathbb{R}[X] \rightarrow \mathbb{R}$ is the linear functional

$$f (= \sum_{\alpha} f_{\alpha} X^{\alpha}) \mapsto L_y(f) = \sum_{\alpha} f_{\alpha} y_{\alpha}.$$

Notice that these conditions are very simple as they require that both moment matrices associated with the sequences y and $\kappa z - y$ are positive semidefinite. Crucial for establishing the result is the generalized multivariate Carleman condition of Nussbaum [9].

2. NOTATION, DEFINITIONS AND PRELIMINARY RESULTS

For a closed set $\mathbf{K} \subset \mathbb{R}^n$, let $\mathcal{B}(\mathbf{K})$ denote its associated Borel σ -algebra, and given a Borel measure μ on $\mathbf{K}$, let $L_\infty(\mathbf{K}, \mu)$ be the standard Banach space of all bounded measurable functions on $\mathbf{K}$ (except possibly on a subset of μ -measure zero).

Let $\mathbb{R}[X]$ be the ring of real polynomials in the variables $X = (X_1, \dots, X_n)$, and let $\Sigma^2 \subset \mathbb{R}[X]$ be its subset of sums of squares. With $\|\cdot\|$ being the euclidean norm of $\mathbb{R}^n$, the notation $\|X\|^2$ stands for the quadratic polynomial $\sum_{i=1}^n X_i^2$. For a real symmetric matrix A , the notation $A \succeq 0$ stands for A is positive semidefinite.

Given a family $\{g_j\}_{j=1}^m \subset \mathbb{R}[X]$, denote by:

- $g_J \in \mathbb{R}[X]$, the polynomial $g_J(X) := \prod_{j \in J} g_j(X)$, for every $J \subseteq \{1, \dots, m\}$, and with the convention $g_\emptyset \equiv 1$.
- $Q(g) \subset \mathbb{R}[X]$, the *quadratic module* generated by the g_j 's, that is,

$$Q(g) := \left\{ u_0 + \sum_{j=1}^m u_j g_j \mid u_j \in \Sigma^2 \quad \forall j = 0, 1, \dots, m \right\}.$$

Given a sequence $y = (y_\alpha)$ indexed in the canonical basis (X^α) of $\mathbb{R}[X]$, let $L_y : \mathbb{R}[X] \rightarrow \mathbb{R}$ be the linear functional

$$f (= \sum_{\alpha \in \mathbb{N}^n} f_\alpha X^\alpha) \mapsto L_y(f) := \sum_{\alpha \in \mathbb{N}^n} f_\alpha y_\alpha.$$

Given a Borel set $\mathbf{K} \subseteq \mathbb{R}^n$, a sequence $y = (y_\alpha)$ has a *representing measure* μ on $\mathbf{K} \subset \mathbb{R}^n$ if

$$y_\alpha = \int_{\mathbf{K}} X^\alpha d\mu, \quad \forall \alpha \in \mathbb{N}^n.$$

A Borel measure μ on $\mathbb{R}^n$ with all its moments finite is said to be *moment determinate* if whenever

$$\int X^\alpha d\mu = \int X^\alpha d\nu, \quad \forall \alpha \in \mathbb{N}^n,$$

holds for some measure ν , then $\nu = \mu$.

When $\mathbf{K} \equiv \mathbb{R}^n$ and if

$$(2) \quad \sum_{k=1}^{\infty} L_y(X_i^{2k})^{-1/2k} = \infty,$$

then by a result from Nussbaum [9], y has a unique representing measure μ ; see e.g. Berg [2]. Condition (2) is called the generalized Carleman condition as an extension to the multivariate case of Carleman's condition for the univariate case.

We next recall several important results concerning necessary and sufficient conditions for a sequence $y = (y_\alpha)$ to have a representing measure on a compact basic semi-algebraic set $\mathbf{K} \subset \mathbb{R}^n$.

So, let $\mathbf{K} \subset \mathbb{R}^n$ be a basic semi-algebraic set defined by:

$$(3) \quad \mathbf{K} := \{x \in \mathbb{R}^n \mid g_j(x) \geq 0, \quad \forall j = 1, \dots, m\},$$

for some polynomials $\{g_j\} \subset \mathbb{R}[X]$.

Theorem 1. *Let $\mathbf{K} \subset \mathbb{R}^n$ be compact and defined as in (3).*

(a) *A given sequence $y = (y_\alpha)$ has a representing measure μ on $\mathbf{K}$ if and only if:*

$$(4) \quad L_y(f^2 g_J) \geq 0 \quad \forall f \in \mathbb{R}[X]; \quad \forall J \subseteq \{1, \dots, m\}.$$

(b) *Assume that for some $M > 0$, the quadratic polynomial $M - \|X\|^2$ is in $Q(g)$. A given sequence $y = (y_\alpha)$ has a representing measure μ on $\mathbf{K}$ if and only if:*

$$(5) \quad L_y(f^2 g_j) \geq 0 \quad \forall f \in \mathbb{R}[X]; \quad \forall j = 0, 1, \dots, m$$

(with the convention $g_0 \equiv 1$).

Theorem 1(a) is the dual part of Schmüdgen's Positivstellensatz [11] whereas its refinement (b) is due to Putinar [10] and Jacobi and Prestel [5].

If one restricts f to have degree less than, say r , then testing conditions (4) and (5) reduces to checking whether some real-valued symmetric matrices are positive semidefinite. Indeed, given $\theta \in \mathbb{R}[X]$, let $M_r(\theta y)$ be the real symmetric matrix with rows and columns indexed in the canonical basis (X^α) of $\mathbb{R}[X]$, and defined by

$$M_r(\theta y)(\alpha, \beta) = L_y(X^{\alpha+\beta} \theta(X)), \quad \forall \alpha, \beta \in \mathbb{N}^n \text{ with } |\alpha|, |\beta| \leq r$$

(where for every $\alpha \in \mathbb{N}^n$, $|\alpha| := \sum_i \alpha_i$). $M_r(\theta y)$ is called the *localizing* matrix associated with y and θ , and when $\theta \equiv 1$, $M_r(y)$ is called the *moment* matrix associated with y .

Then testing condition (4) (resp. (5)) for all $f \in \mathbb{R}[X]$ with $\deg f \leq r$, is equivalent to checking whether $M_r(g_J y) \succeq 0$ for all $J \subseteq \{1, \dots, m\}$ (resp. $M_r(g_j y) \succeq 0$ for all $j = 0, \dots, m$), i.e., computing the smallest eigenvalue of 2^m (resp. $m+1$) real symmetric matrices. If y is taken as unknown, then the LMIs (Linear Matrix Inequalities) $M_r(g_J y) \succeq 0$ (or $M_r(g_j y) \succeq 0$) on y define the set of sequences y which have a representing measure μ on $\mathbf{K}$. Importantly, this set is convex.

Finally, given $\alpha \in \mathbb{N}^n$, let $g^\alpha, (1-g)^\alpha \in \mathbb{R}[X]$ be the polynomials

$$g^\alpha(X) := \prod_{j=1}^m g_j(X)^{\alpha_j}; \quad (1-g)^\alpha(X) := \prod_{j=1}^m (1-g_j(X))^{\alpha_j}.$$

Theorem 2. *Let $\mathbf{K} \subset \mathbb{R}^n$ be as in (3), and assume that the g_j 's are normalized, i.e., $0 \leq g_j \leq 1$ on $\mathbf{K}$ for all $j = 1, \dots, m$, and that the family $(0, 1, \{g_j\})$ generates the algebra $\mathbb{R}[X]$; equivalently $\mathbb{R}[X] = \mathbb{R}[g_1, \dots, g_m]$.*

A given sequence $y = (y_\alpha)$ has a representing measure μ on $\mathbf{K}$ if and only if:

$$(6) \quad L_y(g^\alpha(1-g)^\beta) \geq 0 \quad \forall \alpha, \beta \in \mathbb{N}^n.$$

Theorem 2 is from Vasilescu [15] (also hidden in a previous result of Krivine [7, 8]). Notice that testing whether (6) holds for all $\alpha, \beta \in \mathbb{N}^n$ with $|\alpha|, |\beta| \leq r$, reduces to checking whether finitely many inequalities are satisfied. If again y is taken as unknown, then the *linear* inequalities (6) on y are another representation of the convex set of feasible sequences y which have a representing measure on $\mathbf{K}$.

So this set can be represented either via LMIs through conditions (4) (or (5)), or via linear inequalities through conditions (6).

3. MAIN RESULT

Let μ be a finite Borel measure on a Borel set $\mathbf{K} \subseteq \mathbb{R}^n$. We want to find a set of necessary and/or sufficient conditions on a sequence $y = (y_\alpha)$ to have a representing measure ν on $\mathbf{K}$ that is *uniformly absolutely continuous* with respect to μ , that is, ν is absolutely continuous w.r.t. μ (denoted $\nu \ll \mu$) and its Radon-Nikodym derivative h belongs to $L_\infty(\mathbf{K}, \mu)$. We first concentrate on the case where $\mathbf{K} \subset \mathbb{R}^n$ is basic compact semi-algebraic set as defined in (3), and then consider the non compact case $\mathbf{K} \equiv \mathbb{R}^n$.

3.1. The compact case.

Theorem 3. *Let $\mathbf{K} \subset \mathbb{R}^n$ be compact and defined as in (3). Let $z = (z_\alpha)$ be the moment sequence of a finite Borel measures μ on $\mathbf{K}$.*

(a) *A sequence $y = (y_\alpha)$ has a representing Borel measure ν on $\mathbf{K}$, uniformly absolutely continuous with respect to μ , if and only if there is some scalar κ such that for every $J \subseteq \{1, \dots, m\}$:*

$$(7) \quad 0 \leq L_y(f^2 g_J) \leq \kappa L_z(f^2 g_J) \quad \forall f \in \mathbb{R}[X].$$

(b) *In addition, if the polynomial $N - \|X\|^2$ belongs to the quadratic module $Q(g)$ then one may replace (7) with the weaker condition*

$$(8) \quad 0 \leq L_y(f^2 g_j) \leq \kappa L_z(f^2 g_j) \quad \forall f \in \mathbb{R}[X]; \quad j = 0, \dots, m$$

(with the convention $g_0 \equiv 1$).

(c) *Suppose that the g_j 's are normalized so that*

$$0 \leq g_j \leq 1 \quad \text{on} \quad \mathbf{K}, \quad \forall j = 1, \dots, m,$$

and that the family $(0, 1, \{g_j\})$ generates the algebra $\mathbb{R}[X]$.

A sequence $y = (y_\alpha)$ has a representing Borel measure ν on $\mathbf{K}$, uniformly absolutely continuous with respect to μ , if and only if there is some scalar κ such that for every $\alpha, \beta \in \mathbb{N}^m$:

$$(9) \quad 0 \leq L_y(g^\alpha(1-g)^\beta) \leq \kappa L_z(g^\alpha(1-g)^\beta).$$

Proof. (a). The *only if* part: Let $d\nu = h d\mu$ for some $0 \leq h \in L_\infty(\mathbf{K}, \mu)$, and let $\kappa := \|h\|_\infty$. Observe that $g_J \geq 0$ on $\mathbf{K}$ for all $J \subseteq \{1, \dots, m\}$. Therefore, for every $J \subseteq \{1, \dots, m\}$:

$$\int f^2 g_J d\nu = \int f^2 g_J h d\mu \leq \kappa \int f^2 g_J d\mu, \quad \forall f \in \mathbb{R}[X],$$

and so (7) is satisfied.

The *if part*: Let y and z be such that (7) holds true. Then by Theorem 1(a), y has a representing finite Borel measure ν on $\mathbf{K}$. In addition, let $\gamma = (\gamma_\alpha)$ with $\gamma_\alpha := \kappa z_\alpha - y_\alpha$ for all $\alpha \in \mathbb{N}^n$. From (7), one has

$$L_\gamma(f^2 g_J) \geq 0, \quad \forall f \in \mathbb{R}[X]; \quad \forall J \subseteq \{1, \dots, m\}.$$

By Theorem 1(a) again, γ has a representing finite Borel measure ψ on $\mathbf{K}$. Moreover, from the definition of the sequence γ

$$\int f d(\psi + \nu) = \int f \kappa d\mu, \quad \forall f \in \mathbb{R}[X],$$

and therefore, as measures on compact sets are moment determinate, $\psi + \nu = \kappa \mu$. Hence $\kappa \mu \geq \nu$ which shows that $\nu \ll \mu$. Finally, write $d\nu = h d\mu$ for some $0 \leq h \in L_1(\mathbf{K}, \mu)$. From $\nu \leq \kappa \mu$ one obtains

$$\int_A (h - \kappa) d\mu \leq 0 \quad \forall A \in \mathcal{B}(\mathbf{K}),$$

and so $0 \leq h \leq \kappa$, μ -almost everywhere on $\mathbf{K}$. Equivalently, $\|h\|_\infty \leq \kappa$, the desired result.

(b) The proof is a verbatim copy of (a), except that instead of invoking Theorem 1(a), we now invoke Theorem 1(b), its Putinar's refinement.

(c) The *only if* part: Let $d\nu = h d\mu$ for some $h \in L_\infty(\mathbf{K}, \mu)$, and let $\kappa := \|h\|_\infty$. As $g \geq 0$ and $(1 - g) \geq 0$ on $\mathbf{K}$,

$$\int g^\alpha (1 - g)^\beta d\nu = \int g^\alpha (1 - g)^\beta h d\mu \leq \kappa \int g^\alpha (1 - g)^\beta d\mu,$$

for all $\alpha, \beta \in \mathbb{N}^m$, and so (9) is satisfied.

The *if part*: Let y and z be such that (9) holds true. By Theorem 2, y has a representing Borel measure ν on $\mathbf{K}$. In addition, let $\gamma = (\gamma_\alpha)$ with $\gamma_\alpha := \kappa z_\alpha - y_\alpha$ for all $\alpha \in \mathbb{N}^n$. From (9), the condition (6) holds true with γ in lieu of y , and therefore, by Theorem 2, γ has a representing Borel measure ψ on $\mathbf{K}$. Moreover, from the definition of γ , one has

$$\int f d(\psi + \nu) = \int f \kappa d\mu, \quad \forall f \in \mathbb{R}[X],$$

and the rest of the proof is the same as that of (a). $\square$

Recalling the definition of localizing matrices, notice that checking (7) for all $f \in \mathbb{R}[X]$ with $\deg f \leq r$, reduces to checking whether

$$(10) \quad 0 \preceq M_r(g_J y) \quad \text{and} \quad M_r(g_J y) \preceq \kappa M_r(g_J z) \quad \forall J \subseteq \{1, \dots, m\},$$

whereas checking (8) for all $f \in \mathbb{R}[X]$ with $\deg f \leq r$ reduces to checking whether

$$(11) \quad 0 \preceq M_r(g_j y) \quad \text{and} \quad M_r(g_j y) \preceq \kappa M_r(g_j z) \quad \forall j = 0, \dots, m.$$

While checking the first LMIs in the left of both (10) and (11) is just evaluating the smallest eigenvalue of the 2^m symmetric matrices $M_r(g_J y)$, $J \subseteq \{1, \dots, m\}$ (resp. the $m+1$ symmetric matrices $M_r(g_j y)$, $j = 0, 1, \dots, m$), checking the second LMIs in both (10) and (11) reduces to solving a semi-definite program with only *one* variable κ , and 2^m (resp. $m+1$) LMIs, a convex optimization problem for which efficient public softwares are now available.

On the other hand, checking (9) or all $f \in \mathbb{R}[X]$ with $\deg f \leq r$, reduces to solving a linear program with only one variable also, i.e., checking whether some interval is nonempty.

Theorem 3(c) is the analogue for compact basic semi-algebraic sets of Corollary 3.1 in Knill [6] for the unit box $[0, 1]^n$.

We next obtain a sufficient condition for the non-compact case.

3.2. The non-compact case. Let μ be a finite Borel measure on $\mathbb{R}^n$. We want to find a sufficient condition on a sequence $y = (y_\alpha)$ to be the moment sequence of a measure ν on $\mathbb{R}^n$ that is *uniformly absolutely continuous* with respect to μ .

Theorem 4. *Let $z = (z_\alpha)$ be the moment sequence of a finite Borel measure μ on $\mathbb{R}^n$, and assume that z satisfies the generalized Carleman condition, i.e.,*

$$(12) \quad \sum_{k=1}^{\infty} L_z(X_i^{2k})^{-1/2k} = \infty.$$

Then a sequence $y = (y_\alpha)$ has a representing Borel measure ν on $\mathbb{R}^n$, uniformly absolutely continuous with respect to μ , if there exists a scalar $0 < \kappa$ such that for all $i = 1, \dots, n$:

$$(13) \quad 0 \leq L_y(f^2) \leq \kappa L_z(f^2) \quad \forall f \in \mathbb{R}[X].$$

Proof. For every $i = 1, \dots, n$, (13) with $f^2 = X_i^{2k}$ yields

$$L_y(X_i^{2k})^{-1/2k} \geq \kappa^{-1/2k} L_z(X_i^{2k})^{-1/2k}, \quad \forall k = 0, 1, \dots,$$

and so, using (12), one obtains

$$\sum_{k=1}^{\infty} L_y(X_i^{2k})^{-1/2k} \geq \sum_{k=1}^{\infty} \kappa^{-1/2k} L_z(X_i^{2k})^{-1/2k} = +\infty,$$

for every $i = 1, \dots, n$, i.e., the generalized Carleman condition (2) holds for the sequence y . Combining this with the first inequality in (13) yields that y has a unique representing Borel measure ν on $\mathbb{R}^n$. It remains to prove that $\nu \ll \mu$ and its density h is in $L_\infty(\mathbb{R}^n, \mu)$.

Let $\gamma = (\gamma_\alpha)$ with $\gamma_\alpha := \kappa z_\alpha - y_\alpha$ for all $\alpha \in \mathbb{N}^n$. Then the second inequality in (13) yields

$$(14) \quad L_\gamma(f^2) \geq 0 \quad \forall f \in \mathbb{R}[X].$$

Next, observe that from (13), for every $i = 1, \dots, n$, and every $k = 0, 1, \dots$,

$$L_\gamma(X_i^{2k}) \leq \kappa L_z(X_i^{2k}),$$

which implies

$$(15) \quad L_\gamma(X_i^{2k})^{-1/2k} \geq \kappa^{-1/2k} L_z(X_i^{2k})^{-1/2k},$$

and so, for every $i = 1, \dots, n$,

$$\sum_{k=1}^{\infty} L_\gamma(X_i^{2k})^{-1/2k} \geq \sum_{k=1}^{\infty} \kappa^{-1/2k} L_z(X_i^{2k})^{-1/2k} = +\infty,$$

i.e., γ satisfies the generalized Carleman condition. In view of (14), γ has a (unique) representing measure ψ on $\mathbb{R}^n$. Next, from the definition of γ , one has

$$\int f d(\psi + \nu) = \kappa \int f d\mu, \quad \forall f \in \mathbb{R}[X].$$

But as μ (and so $\kappa\mu$) satisfies Carleman condition (12), $\kappa\mu$ is moment determinate and therefore, $\kappa\mu = \nu + \psi$.

Hence $\nu \ll \mu$ follows from $\nu \leq \kappa\mu$. Finally, writing $d\nu = h d\mu$ for some nonnegative $h \in L_1(\mathbb{R}^n, \mu)$, and using $\nu \leq \kappa\mu$, one obtains

$$\int_A (h - \kappa) d\mu \leq 0 \quad \forall A \in \mathcal{B}(\mathbb{R}^n),$$

and so $0 \leq h \leq \kappa$, μ -almost everywhere on $\mathbb{R}^n$. Equivalently, $\|h\|_\infty \leq \kappa$, the desired result. $\square$

Observe that condition (13) is extremely simple as it is equivalent to stating that

$$\kappa M_r(z) \succeq M_r(y) \succeq 0, \quad \forall r = 0, 1, \dots$$

where $M_r(z)$ (resp. $M_r(y)$) is the r -moment matrix associated with the sequence z (resp. y). Finally, a sufficient condition for (12) to hold is

$$\int \exp |x_i| d\mu < \infty, \quad \forall i = 1, \dots, n.$$

ACKNOWLEDGEMENTS

The author acknowledges financial support by the (french) National Research Agency (ANR), under grant NT05-3-41612.

REFERENCES

- [1] AKHIEZER, N.I. (1965). *The classical moment problem and some related questions in analysis*, Hafner Publishing Company, New York.
- [2] BERG, C. (1980). The multidimensional moment problem and semi-groups, in: *Moment in Mathematics*, ed. H.J. Landau, Amer. Math. Soc., Providence, RI., pp. 110–124.
- [3] CURTO, R.E. AND FIALKOW, L.A. (2000). The truncated complex K -moment problem, *Trans. Amer. Math. Soc.* **352**, 2825–2855.
- [4] HANDELMAN, D. (1988). Representing polynomials by positive linear functions on compact convex polyhedra, *Pac. J. Math.* **132**, 35–62.
- [5] JACOBI, T. AND PRESTEL, A. (2001). Distinguished representations of strictly positive polynomials, *J. reine angew. Math.* **532**, 223–235.
- [6] KNILL, O. (2000). On Hausdorff’s moment conditions in higher dimensions, Preprint, Department. of Mathematics, University of Arizona, Tucson, AZ 85721, USA. Available at www.math.harvard.edu/~knill/
- [7] KRIVINE, J.L. (1964). Anneaux préordonnés, *J. Anal. Math.* **12**, 307–326.
- [8] KRIVINE, J.L. (1964). Quelques propriétés des préordres dans les anneaux commutatifs unitaires, *C.R. Acad. Sci. Paris* **258**, 3417–3418.
- [9] NUSSBAUM, A.E. (1966). Quasi-analytic vectors, *Ark. Mat.* **6**, 179–191.
- [10] PUTINAR, M. (1993). Positive polynomials on compact semi-algebraic sets, *Ind. Univ. Math. J.* **42**, 969–984.
- [11] SCHMÜDGEN, K. (1991). The K -moment problem for compact semi-algebraic sets, *Math. Ann.* **289**, 203–206.
- [12] SHOHAT, J.A. AND TAMARKIN, J.D. (1943). *The problem of moments*, Mathematical Surveys **1**, Amer. Math. Soc. Providence, RI.
- [13] SIMON, B. (1998). The classical moment problem as a self-adjoint finite difference operator, *Adv. Math.* **137**, 82–203.
- [14] VANDENBERGHE, L. AND BOYD, S. (1996). Semidefinite programming, *SIAM Review* **38**, 49–95.
- [15] VASILESCU, F.-H. (2003). Spectral measures and moment problems, in: *Spectral Theory and Its Applications*, Theta Foundation, Bucharest, Romania, pp. 173–215.

LAAS-CNRS AND INSTITUTE OF MATHEMATICS, LAAS 7 AVENUE DU COLONEL ROCHE, 31077 TOULOUSE CÉDEX 4 (FRANCE), URL: WWW.LAAS.FR/~LASSERRE
E-mail address: `lasserre@laas.fr`